

A discrete version of the inverse scattering problem and the J-matrix method

S. A. Zaytsev¹

*Department of Physics, Khabarovsk State University of Technology,
Tikhookeanskaya 136, Khabarovsk 680035, Russia*

Abstract

The problem of the Hamiltonian matrix in the oscillator and orthogonalized Laguerre basis construction from a given S-matrix is treated in the context of the algebraic analogue of the Marchenko method.

1 Introduction

The J-matrix [1] theory of scattering is based on the fact that the ℓ th partial wave kinetic energy or the Coulomb Hamiltonian H^0 is represented in a certain square-integrable basis set by an infinite symmetric tridiagonal matrix. In the harmonic oscillator and the Laguerre basis sets $\{\phi_n^\ell\}_{n=0}^\infty$ the eigenvalue problem for H^0 can be solved analytically. The J-matrix method yields an exact solution to a model scattering Hamiltonian where the given short-range potential is approximated by truncating in a finite subset $\{\phi_n^\ell\}_{n=0}^{N-1}$.

In Refs. [2, 3] an inverse scattering formalism within the J-matrix method has been proposed, where the matrix $\|V_{n,m}\|$ of the potential

$$V_\ell(r, r') = \hbar\omega \sum_{n,m=0}^{N-1} \phi_n^\ell(x) V_{n,m} \phi_m^\ell(x') \quad (1)$$

with the oscillator form factors

$$\phi_n^\ell(x) = (-1)^n \sqrt{\frac{2n!}{\rho\Gamma(n+\ell+\frac{3}{2})}} x^{\ell+1} e^{-x^2/2} L_n^{\ell+1/2}(x^2) \quad (2)$$

is determined from a given S-matrix. Here, $x = r/\rho$ is the relative coordinate in units of the oscillator radius $\rho = \sqrt{\hbar/\mu\omega}$, μ is the reduced mass.

Obviously a correlation can be made between the J-matrix method and a discrete model of quantum mechanics, within of which a finite-difference Schrödinger equation is used. As a result the J-matrix versions of the

¹This work has been done partially while the author was visiting the Institute for Nuclear Theory, University of Washington.

Gel'fand-Levitan-Marchenko method algebraic analogue can be formulated. For instance, the J-matrix method formally and computationally is quite similar to the R-matrix theory. It is this analogy that the previous J-matrix version of inverse scattering theory [2, 3, 4] [also see [5]] leans upon. Within the J-matrix approach the discrete representation of the Green function in finite subspace of the basis functions $\{\phi_n^\ell\}_{n=0}^{N-1}$

$$\mathbf{G}(\epsilon) = (\epsilon \mathbf{I} - \mathbf{h})^{-1} \quad (3)$$

is used. Here, $\mathbf{I}$ is identity matrix and $\mathbf{h}$ is the truncated Hamiltonian matrix of order N in the oscillator basis (2). We measure the energy E in the units of the oscillator basis parameter $\hbar\omega$, i.e. $E = \hbar\omega\epsilon$ and $\epsilon = q^2/2$, where q is the dimensionless momentum: $q = k\rho$. In particular, the element $[\mathbf{G}(\epsilon)]_{N-1, N-1} \equiv \mathcal{P}_N(\epsilon)$ can be presented in the two rational forms [2, 6]:

$$\mathcal{P}_N(\epsilon) = \frac{\prod_{j=0}^{N-2} (\epsilon - \mu_j)}{\prod_{j=0}^{N-1} (\epsilon - \lambda_j)} \quad (4)$$

where $\{\lambda_j\}_{j=0}^{N-1}$ and $\{\mu_j\}_{j=0}^{N-2}$ satisfy the interlacing property

$$\lambda_0 < \mu_0 < \lambda_1 < \dots < \lambda_{N-2} < \mu_{N-2} < \lambda_{N-1},$$

and [7]

$$\mathcal{P}_N(\epsilon) = \sum_{j=0}^{N-1} \frac{Z_{N-1,j}^2}{\epsilon - \lambda_j}. \quad (5)$$

Here, $\{Z_{N-1,j}\}_{j=0}^{N-1}$ are the elements of the N th row of the eigenvector matrix $\mathbf{Z}$ of the truncated Hamiltonian matrix $\mathbf{h}$. The sets $\{\lambda_j\}_{j=0}^{N-1}$, $\{\mu_j\}_{j=0}^{N-2}$ and $\{\lambda_j\}_{j=0}^{N-1}$, $\{Z_{N-1,j}\}_{j=0}^{N-1}$ are derived from the S-matrix which is intimately connected with $\mathcal{P}_N$ [see Eq. (39)]. Notice that the both sets of the spectral parameters determine unique [apart from the off diagonal elements sign] Hamiltonian matrix $\mathbf{h}$ of a Jacobi form [8, 9]

$$\|h_{n,m}\| = \begin{pmatrix} a_0 & b_1 & & & \\ b_1 & a_1 & b_2 & & 0 \\ & b_2 & a_2 & b_3 & \\ & & b_3 & \times & \times \\ 0 & & & \times & \times & b_{N-1} \\ & & & & b_{N-1} & a_{N-1} \end{pmatrix}. \quad (6)$$

Hence, the sought-for potential matrix $\|V_{n,m}\|$ is also of a Jacobi form. Recall that the kinetic energy operator

$$H^0 = \frac{\hbar^2}{2\mu} \left(-\frac{d^2}{dr^2} + \frac{\ell(\ell+1)}{r^2} \right) \quad (7)$$

matrix representation $\|T_{n,m}^\ell\| = \frac{1}{\hbar\omega} \|H_{n,m}^0\|$ in the harmonic oscillator basis (2) is symmetric tridiagonal [1, 7]:

$$\begin{aligned} T_{n,n}^\ell &= \frac{1}{2} \left(2n + \ell + \frac{3}{2} \right), \\ T_{n,n+1}^\ell &= T_{n+1,n}^\ell = -\frac{1}{2} \sqrt{(n+1) \left(n + \ell + \frac{3}{2} \right)}. \end{aligned} \quad (8)$$

Thus the inverse scattering problem within the J-matrix approach admits of the solution in the tridiagonal Hamiltonian matrix form. In this regard the J-matrix method is similar to a discrete model of quantum mechanics, in the framework of which the Hamiltonian matrix representation is also symmetric tridiagonal. Note that the tridiagonal matrix representation of both the kinetic energy operator and the Hamiltonian is fundamental for a finite-difference analogue of the Gel'fand-Levitan equations [see e.g. [10]] as well as for a discrete version [11] of the Marchenko equations. As shown below, the similarity between the J-matrix method and a finite-difference approach can be plainly extended to the inverse scattering formalism as well. In the present paper, in particular, an inverse scattering J-matrix approach via the Marchenko equations (JME) is given.

In JME the expansion coefficients c_n of the wave function ψ in terms of the L^2 basis set play a role similar to that of the values $\psi_n = \psi(x_n)$ at points $x_n = n\Delta$ within the finite-difference inverse scattering approach. Here, the completeness relation for the solutions c_n of the Schrödinger equation discrete analogue is also exploited which involves an integration $c_n \overline{c_m}$ over the energy from zero to infinity. This raises the question as to whether the integrals converge. As shown below, taking account of the phase shift $\delta_\ell^N(k)$ corresponding to the potential (1) of finite rank N asymptotic behavior at large k provides the convergence of the integrals. Generally, with a potential (1) matrix of finite order N it is possible to reproduce the phase shift δ_ℓ only on a finite energy interval $[0, \epsilon_0]$ with $\epsilon_0 < \lambda_{N-1}$. This is why the eigenvalue λ_{N-1} and the corresponding eigenvector component $Z_{N-1, N-1}$ are the variational parameters within the previous J-matrix version [2, 3, 4, 5] of the inverse scattering theory. By contrast, in JME the phase shift, even if modified in accordance with the δ_ℓ^N asymptotic feature, on infinite energy interval is used. As a result JME has not variational parameters (apart from N and ρ).

The elements of the J-matrix method formalism are presented in Sec. 2. In Sec. 3 the inverse scattering J-matrix approach in the context of the Marchenko equations is formulated. The features of JME numerical realization are discussed in Sec. 4. In Sec. 5 JME is expanded to the Laguerre basis case. Here, we are dealing with the tridiagonal Hamiltonian matrix construction in an orthogonalized Laguerre basis set, in which the kinetic energy operator matrix is also tridiagonal. In Sec. 6 we summarize our conclusions.

2 The direct problem

The oscillator-basis J-matrix formalism is discussed in detail elsewhere. We present here only some relations needed for understanding the inverse scattering J-matrix approach. Within the J-matrix method, the radial wave function $\psi(k, r)$ is expanded in an oscillator function (2) series

$$\psi(k, r) = \sum_{n=0}^{\infty} c_n(k) \phi_n^\ell(x). \quad (9)$$

In the assumption that the Hamiltonian matrix is of the form (6) the functions c_n are the solutions to the set of equations

$$\begin{aligned} a_0 c_0(k) + b_1 c_1(k) &= \epsilon c_0(k) \\ b_n c_{n-1}(k) + a_n c_n(k) + b_{n+1} c_{n+1}(k) &= \epsilon c_n(k), \quad n = 1, 2, \dots \end{aligned} \quad (10)$$

The asymptotic behavior of $c_n(k)$ for $k > 0$ as $n \rightarrow \infty$ is given by

$$c_n(k) = f_n(k) \equiv \frac{i}{2} \left[\mathcal{C}_{n,\ell}^{(-)}(q) - S(k) \mathcal{C}_{n,\ell}^{(+)}(q) \right]. \quad (11)$$

Here, the functions

$$\begin{aligned} \mathcal{C}_{n,\ell}^{(\pm)}(q) &= \mathcal{C}_{n,\ell}(q) \pm i \mathcal{S}_{n,\ell}(q), \\ \mathcal{S}_{n,\ell}(q) &= \sqrt{\frac{\pi \rho n!}{\Gamma(n+\ell+\frac{3}{2})}} q^{\ell+1} e^{-q^2/2} L_n^{\ell+1/2}(q^2), \\ \mathcal{C}_{n,\ell}(q) &= \sqrt{\frac{\pi \rho n!}{\Gamma(n+\ell+\frac{3}{2})}} \frac{\Gamma(\ell+1/2)}{\pi q^\ell} e^{-q^2/2} F(-n-\ell-1/2, -\ell+1/2; q^2) \end{aligned} \quad (12)$$

obey the “free” equations

$$T_{n,n-1}^\ell e_{n-1}(q) + T_{n,n}^\ell e_n(q) + T_{n,n+1}^\ell e_{n+1}(q) = \epsilon d_n(q), \quad n = 1, 2, \dots \quad (13)$$

$\mathcal{S}_{n,\ell}$ satisfy in addition the equation

$$T_{0,0}^\ell \mathcal{S}_{0,\ell}(q) + T_{0,1}^\ell \mathcal{S}_{1,\ell}(q) = \epsilon \mathcal{S}_{0,\ell}(q). \quad (14)$$

Besides, $\mathcal{S}_{n,\ell}$ meet the completeness relation

$$\frac{2}{\pi} \int_0^\infty dk \mathcal{S}_{n,\ell}(q) \mathcal{S}_{m,\ell}(q) = \delta_{n,m}. \quad (15)$$

Notice that

$$\begin{aligned} \tilde{S}(r) &= \sum_{n=0}^\infty \mathcal{S}_{n,\ell}(q) \phi_n^\ell(x), \\ \tilde{C}(r) &= \sum_{n=0}^\infty \mathcal{C}_{n,\ell}(q) \phi_n^\ell(x), \end{aligned}$$

subject to the asymptotic condition [7]

$$\begin{aligned} \tilde{S}(r) &\underset{r \rightarrow \infty}{\sim} \sin(kr - \ell\pi/2), \\ \tilde{C}(r) &\underset{r \rightarrow \infty}{\sim} \cos(kr - \ell\pi/2). \end{aligned} \quad (16)$$

As for the coefficients of the expansion

$$\psi_\nu(r) = \sum_{n=0}^\infty c_n(i\kappa_\nu) \phi_n^\ell(x) \quad (17)$$

of the normalized bound state wave function ψ_ν with the energy $-\kappa_\nu^2/2$,

$$c_n(i\kappa_\nu) = f_n(i\kappa_\nu) \equiv \mathcal{M}_\nu i^\ell \mathcal{C}_{n,\ell}^{(+)}(i\kappa_\nu \rho) \quad (18)$$

holds as $n \rightarrow \infty$. Here, $\mathcal{M}_\nu$ is the bound state normalization constant which is related to the residue of the S-matrix [13]:

$$i \operatorname{Res}_{k=i\kappa_\nu} S(k) = (-1)^\ell \mathcal{M}_\nu^2. \quad (19)$$

It can be easy verified that from the completeness relation for the solutions $\psi(k, r)$, $\psi_\nu(r)$ [14]

$$\frac{2}{\pi} \int_0^\infty dk \psi(k, x) \overline{\psi(k, y)} + \sum_\nu \psi_\nu(x) \overline{\psi_\nu(y)} = \delta(x - y) \quad (20)$$

it follows that

$$\frac{2}{\pi} \int_0^\infty dk c_n(k) \overline{c_m(k)} + \sum_\nu c_n(i\kappa_\nu) \overline{c_m(i\kappa_\nu)} = \delta_{n,m}. \quad (21)$$

3 The inverse problem

To take advantage of the algebraic analogue of the Marchenko method it is essential that there exist coefficients $K_{n,m}$ [independent of k] such that

$$c_n(k) = \sum_{m=n}^{\infty} K_{n,m} f_m(k). \quad (22)$$

By analogy with Ref. [11] assume that

$$c_n(k) = f_n(k), \quad n \geq N \quad (23)$$

[N specifies the order of a potential matrix]. If f_N and f_{N+1} are inserted [instead of respectively c_N and c_{N+1}] into Eq. (10) for $n = N$, we obtain, in view of Eq. (13),

$$\begin{aligned} c_{N-1}(k) = & (T_{N,N-1}^{\ell} f_{N-1}(k) + [T_{N,N}^{\ell} - a_N] f_N(k) + \\ & + [T_{N,N+1}^{\ell} - b_{N+1}] f_{N+1}(k)) / b_N. \end{aligned} \quad (24)$$

Then, using the three-term recursion relation (13) with every $n = N - 1, \dots, 1$ we obtain

$$c_n(k) = \sum_{m=n}^{2N-n-1} K_{n,m} f_m(k), \quad n = 0, 1, \dots, N-1 \quad (25)$$

[which in the limit $N \rightarrow \infty$ gives (22)].

The coefficients $K_{n,m}$ are found from the completeness relation (21). From the condition of the orthogonality of c_n and every c_m , $m > n$ follows the condition of the orthogonality of c_n and every f_m , $m > n$, i. e.

$$\frac{2}{\pi} \int_0^{\infty} dk c_n(k) \overline{f_m(k)} + \sum_{\nu} c_n(i\kappa_{\nu}) \overline{f_m(i\kappa_{\nu})} = 0, \quad m > n. \quad (26)$$

Inserting the expansion of (25) in (26) gives the system of linear equations in $K_{n,m}$

$$K_{n,n} Q_{n,m} + \sum_{p=n+1}^{2N-n-1} K_{n,p} Q_{p,m} = 0, \quad m > n. \quad (27)$$

Then, inserting Eq. (25) into (21) and putting $n = m$, we obtain, in view of Eq. (26), the equation in $K_{n,n}$

$$K_{n,n} \left(K_{n,n} Q_{n,n} + \sum_{p=n+1}^{2N-n-1} K_{n,p} Q_{p,n} \right) = 1. \quad (28)$$

Note that from Eq. (27) it follows that $K_{n,m}$, $m > n$ are proportional to $K_{n,n}$. In Eqs. (27) and (28) $Q_{n,m}$ are defined from the scattering data by

$$Q_{n,m} = \frac{2}{\pi} \int_0^\infty dk f_n(k) \overline{f_m(k)} + \sum_\nu f_n(i\kappa_\nu) \overline{f_m(i\kappa_\nu)}. \quad (29)$$

The elements a_n and b_n of the sought-for Hamiltonian matrix (6) are related to $K_{n,m}$ by the equations

$$\begin{aligned} a_n &= T_{n,n}^\ell + \frac{K_{n,n+1}}{K_{n,n}} T_{n+1,n}^\ell - \frac{K_{n-1,n}}{K_{n-1,n-1}} T_{n,n-1}^\ell, \\ b_n &= \frac{K_{n,n}}{K_{n-1,n-1}} T_{n,n-1}^\ell, \quad n = 1, 2, 3, \dots \end{aligned} \quad (30)$$

a_0 is specified by the solutions c_0 and c_1 to Eq. (10).

4 A numerical realization

To this point the assumption has been made that the phase shift $\delta_\ell(k)$ is a continuous function of the wave number k that meets the conditions [14]

$$\delta_\ell(\infty) = 0, \quad \int_0^\infty k^{-1} |\delta_\ell(k)| dr < \infty.$$

In this case δ_ℓ must satisfy stringent requirements. Indeed, the integrated function in r.h.s. of Eq.(29) can be expressed in the form of a product $g_n g_m$ of (real) functions

$$g_n(q) = \mathcal{S}_{n\ell}(q) \cos \delta_\ell + \mathcal{C}_{n\ell}(q) \sin \delta_\ell. \quad (31)$$

It is obviously that a sufficient condition to the convergence of the integrals in (29) is that functions g_n are square-integrable. Notice that from (12) follows

$$\mathcal{S}_{n\ell}(q) \underset{q \rightarrow \infty}{\sim} (-1)^n \sqrt{\frac{\pi \rho}{n! \Gamma(n + \ell + \frac{3}{2})}} q^{2n+\ell+1} e^{-q^2/2}, \quad (32)$$

i.e. the first term in (31) decays exponentially at asymptotically large q . However, $\mathcal{C}_{n\ell}$ grows exponentially with increasing q :

$$\mathcal{C}_{n\ell}(q) \underset{q \rightarrow \infty}{\sim} (-1)^{n+1} \frac{\sqrt{\pi \rho n! \Gamma(n + \ell + \frac{3}{2})}}{\pi} q^{-(2n+\ell+2)} e^{q^2/2}. \quad (33)$$

This suggests that the phase shift δ_ℓ must decay rapidly enough to provide the convergence of the integral in r.h.s. of Eq. (29).

Actually the phase shift δ_ℓ^N corresponding to the potential (1) of rank N [7]

$$\tan \delta_\ell^N = -\frac{\mathcal{S}_{N-1,\ell}(q) - \mathcal{P}_N(\epsilon) T_{N-1,N}^\ell \mathcal{S}_{N,\ell}(q)}{\mathcal{C}_{N-1,\ell}(q) - \mathcal{P}_N(\epsilon) T_{N-1,N}^\ell \mathcal{C}_{N,\ell}(q)}, \quad (34)$$

as seen in Eqs. (32), (33), fulfills even more strict requirement

$$\delta_\ell^N \underset{q \rightarrow \infty}{\sim} \frac{\pi (2N + \ell - \frac{1}{2})}{(N-1)! \Gamma(N + \ell + \frac{1}{2})} q^{4N+2\ell-3} e^{-q^2}. \quad (35)$$

Because of the restriction (35) on the phase shift δ_ℓ^N the potential (1) of finite rank N generally is incapable to describe the scattering data on the infinite interval $k \in [0, \infty)$. At most, we can set ourselves the task of constructing the potential (1) that describes the experimental phase shift δ_ℓ on some finite interval $[0, k_0]$, since generally δ_ℓ needs to be modified in the region $k > k_0$ to provide at least the convergence of the integrals in Eq. (29).

As an example we consider the s -wave scattering case. The “experimental” phase shift δ_ℓ [dotted curve in figure 1] is that of the scattering on the potential given by straight well with the depth V_0 : $\sqrt{\frac{2\mu}{\hbar^2}} V_0 R = 1.5$. The potential (1) is sought for that describes the phase shift on the interval $[0, k_0]$, $k_0 R = 6$ [in figure 1 crosses represent a modified phase shift]. The phase shift δ_ℓ^N corresponding to the resulting potential (1) of rank $N = 6$ and $\rho = \frac{1}{2}R$ is shown in figure 1 [solid curve]. Notice that $\mathcal{C}_{n\ell}$ explodes exponentially with increasing q . Thus, the contribution from the region $q > q_0 = \rho k_0 = 3$ to the integral in Eq. (29) may become overwhelming [see figure 2 where $g_0(q)^2$ is plotted], with the result that the method fails. Matters can be improved by a transition to lesser ρ that shifts q_0 to a region where $\mathcal{C}_{n\ell}$ is not that large, or replacing δ_ℓ at $q > q_0$ with a function that decays rapidly enough.

In the second example, a scattering data on the potential with a bound state has been used as input. The phase shift δ_ℓ [dotted curve in figure 3] corresponds to the s -wave scattering on a spherically symmetric potential in the form of straight well. The well parameter $\sqrt{\frac{2\mu}{\hbar^2}} V_0 R = 2$ determines the bound state with the energy $E = -\kappa^2$, $\kappa R = 0.638045$ and the asymptotic normalization constant $\mathcal{M}R^{1/2} = 1.583324$. k_0 and ρ have been taken the same as in the first example [the modified phase shift is represented by crosses in figure 3]. It is well known that a phase shift does not depend on energy positions and asymptotic normalization constants of bound states. Thus, the inverse scattering problem in the presence of a bound state can be split into two steps.

On the first step we focus on the describing the phase shift and, in spite of the whole of scattering data is used [see Eq.(29)], do not seek to describe the

bound state with high degree of accuracy. The phase shift δ_ℓ^N corresponding to the potential (1) parameters, which together with κR and $\mathcal{M}R^{1/2}$ are presented in the left half of Table, is shown in figure 3 [solid curve].

On the second step, to improve the description of the bound states we use the relationship (19) between the poles and residues of the S-matrix and the characteristics of the bound states [see e.g. Ref. [5]]. Here, the smallest eigenvalue λ_0 and the corresponding eigenvector component $Z_{N-1,0}$ associated with the bound state are found from the system

$$\sum_{j=0}^{N-1} Z_{N-1,j}^2 = 1, \quad (36)$$

$$\mathcal{P}_N(-\kappa^2 \rho^2 / 2) = \frac{1}{T_{N-1,N}^\ell} \frac{\mathcal{C}_{N-1,\ell}^{(+)}(i\kappa\rho)}{\mathcal{C}_{N,\ell}^{(+)}(i\kappa\rho)}, \quad (37)$$

$$\frac{\mathcal{C}_{N-1,\ell}^{(-)}(i\kappa\rho) - \mathcal{P}_N(-\kappa^2 \rho^2 / 2) T_{N-1,N}^\ell \mathcal{C}_{N,\ell}^{(-)}(i\kappa\rho)}{\frac{d}{dq} \left\{ \mathcal{C}_{N-1,\ell}^{(+)}(q) - \mathcal{P}_N(q^2 / 2) T_{N-1,N}^\ell \mathcal{C}_{N,\ell}^{(+)}(q) \right\} \Big|_{q=i\kappa\rho}} = i(-1)^{\ell+1} \rho \mathcal{M}^2. \quad (38)$$

Eqs. (37), (38) are derived from the S-matrix formula for the potential (1) [7]

$$S_\ell^N = \frac{\mathcal{C}_{N-1,\ell}^{(-)}(q) - \mathcal{P}_N(\epsilon) T_{N-1,N}^\ell \mathcal{C}_{N,\ell}^{(-)}(q)}{\mathcal{C}_{N-1,\ell}^{(+)}(q) - \mathcal{P}_N(\epsilon) T_{N-1,N}^\ell \mathcal{C}_{N,\ell}^{(+)}(q)} \quad (39)$$

and Eq. (19). Notice that the component $Z_{N-1,N-1}$ corresponding to the leading eigenvalue λ_{N-1} is involved to meet the normalization condition (36). The phase shift is scarcely affected by changing the parameters $\{\lambda_0, Z_{N-1,0}, Z_{N-1,N-1}\}$ from the initial values obtained on the first step to the ones that are evaluated from Eqs.(36) - (38). The potential parameters, which provide the correct values of κR and $\mathcal{M}R^{1/2}$, are presented in the right half of Table.

5 The Laguerre basis

5.1 Preliminaries

For simplicity's sake we restrict our consideration to the scattering of neutral particles. However, the resulting equations still stand in the presence of the repulsive Coulomb interaction. The potential sought is given by the expression

$$V_\ell(r, r') = \frac{\hbar^2}{2\mu} \sum_{n,m=0}^{N-1} \bar{\phi}_n^\ell(x) V_{n,m} \bar{\phi}_m^\ell(x') \quad (40)$$

where the functions $\bar{\phi}_n^\ell$

$$\bar{\phi}_n^\ell(x) = \frac{n!}{r(n+2\ell+1)!} \phi_n^\ell(x) \quad (41)$$

are bi-orthogonal to the base Laguerre functions ϕ_n^ℓ :

$$\phi_n^\ell(x) = (2br)^{\ell+1} e^{-br} L_n^{2\ell+1}(2br), \quad (42)$$

i.e.

$$\int_0^\infty dr \bar{\phi}_n^\ell(x) \phi_m^\ell(x) = \delta_{n,m}. \quad (43)$$

Here, b is the scale parameter: $x = br$.

The coefficients u_n of the expansion

$$\psi(k, r) = \sum_{n=0}^\infty u_n(k) \phi_n^\ell(x) \quad (44)$$

of the Schrödinger equation regular solution $\psi(k, r)$ satisfy the system of equations

$$(h_{n,m}^0 + V_{n,m}) u_m(k) = k^2 A_{n,m}^\ell u_m(k), \quad n = 0, 1, \dots \quad (45)$$

Here, $\|h_{n,m}^0\|$ is the symmetric tridiagonal matrix of the reference Hamiltonian $\frac{2\mu}{\hbar^2} H^0$ (7) calculated in the basis (42) [1]:

$$\begin{aligned} h_{n,n}^0 &= b \frac{(n+2\ell+1)!}{n!} (n + \ell + 1), \\ h_{n,n+1}^0 &= h_{n+1,n}^0 = b \frac{(n+2\ell+2)!}{2n!}, \quad n = 0, 1, \dots \end{aligned} \quad (46)$$

$\|A_{n,m}^\ell\|$ signifies the basis-overlap matrix

$$A_{n,m}^\ell = \int_0^\infty dr \phi_n^\ell(x) \phi_m^\ell(x) \quad (47)$$

which is also of Jacobi form:

$$\begin{aligned} A_{n,n}^\ell &= \frac{(n+2\ell+1)!}{bn!} (n + \ell + 1), \\ A_{n,n+1}^\ell &= A_{n+1,n}^\ell = -\frac{(n+2\ell+2)!}{2bn!}, \quad n = 0, 1, \dots \end{aligned} \quad (48)$$

The asymptotic behaviour of the coefficients $u_n(k)$, $k > 0$, as $n \rightarrow \infty$ is given by the following expression:

$$u_n(k) = w_n(k) \equiv \frac{i}{2} [\mathcal{C}_{n,\ell}^{(-)}(k) - S(k) \mathcal{C}_{n,\ell}^{(+)}(k)] \quad (49)$$

where the functions [7]

$$\begin{aligned} \mathcal{C}_{n,\ell}^{(\pm)}(k) &= -\frac{n!}{(n+\ell+1)!} \frac{(-\xi)^{\pm(n+1)}}{(2 \sin \zeta)^\ell} {}_2F_1(-\ell, n+1; n+\ell+2; \xi^{\pm 2}), \\ \xi &= e^{i\zeta} = \frac{ib-k}{ib+k}, \end{aligned} \quad (50)$$

obey the inhomogeneous “free” equation

$$J_{n,m}^\ell(k) \mathcal{C}_{m,\ell}^{(\pm)}(k) = \delta_{n,0} \frac{k}{\mathcal{S}_{0,\ell}(k)}, \quad n = 0, 1, \dots \quad (51)$$

Here, $\|J_{n,m}^\ell(k)\| = \|h_{n,m}^0 - k^2 A_{n,m}^\ell\|$ is the so-called J-matrix. $\mathcal{S}_{n,\ell}$ are the solutions of the system of equations

$$J_{n,m}^\ell(k) \mathcal{S}_{m,\ell}(k) = 0, \quad n = 0, 1, \dots, \quad (52)$$

$$\mathcal{S}_{n,\ell}(k) = \frac{\ell! (2 \sin \zeta)^{\ell+1}}{2(2\ell+1)!} (-\xi)^n {}_2F_1(-n, \ell+1; 2\ell+2; 1-\xi^{-2}). \quad (53)$$

It can easily be shown that the completeness relation for the functions $\mathcal{S}_{n,\ell}$ of Ref. [15] can be rewritten as

$$\frac{2}{\pi} \int_0^\infty dk \mathcal{S}_{n,\ell}(k) A_{n',m}^\ell \mathcal{S}_{m,\ell}(k) = \delta_{n,n'}. \quad (54)$$

The coefficients $u_n(i\kappa_\nu)$ of the expansion of the bound state normalized wave function $\psi_\nu(r)$ with the energy $-\kappa_\nu^2$ have the following asymptotic behaviour

$$u_n(i\kappa_\nu) = w_n(i\kappa_\nu) \equiv \mathcal{M}_\nu i^\ell \mathcal{C}_{n,\ell}^{(+)}(i\kappa_\nu) \quad (55)$$

as $n \rightarrow \infty$.

Notice that the sine-like J-matrix solutions $\tilde{S}(r) = \sum_{n=0}^\infty \mathcal{S}_{n,\ell}(k) \phi_n^\ell(x)$ and the cosine-like one $\tilde{C}(r) = \sum_{n=0}^\infty \mathcal{C}_{n,\ell}(k) \phi_n^\ell(x)$, where $\mathcal{C}_{n,\ell}(k) = \frac{1}{2} (\mathcal{C}_{n,\ell}^{(+)}(k) + \mathcal{C}_{n,\ell}^{(-)}(k))$, have the asymptotic behaviour (16).

The the completeness relation (20) is transformed into

$$\frac{2}{\pi} \int_0^\infty u_n(k) A_{n',m}^\ell \overline{u_m(k)} + \sum_\nu u_n(i\kappa_\nu) A_{n',m}^\ell \overline{u_m(i\kappa_\nu)} = \delta_{n,n'}. \quad (56)$$

5.2 Inverse problem

In the framework of the J-matrix version [4] of the inverse scattering problem the spectral parameter set $\{\lambda_j, Z_{N-1,j}\}_{j=0}^{N-1}$ is obtained from the scattering data of the truncated Hamiltonian matrix of order N in the orthogonal basis $\varphi_n^\ell = \sum_{m=0}^{N-1} D_{n,m}^\ell \phi_m^\ell$, where

$$D_{n,m}^\ell = \begin{cases} d_n^\ell, & n \geq m, \\ 0, & n < m, \end{cases} \quad d_n^\ell = \sqrt{\frac{2b n!}{(n+2\ell+2)!}}, \quad (57)$$

i. e.

$$\varphi_n^\ell(x) = d_n^\ell (2br)^{\ell+1} e^{-br} L_n^{2\ell+2}(2br). \quad (58)$$

Clearly the set $\{\lambda_j, Z_{N-1,j}\}_{j=0}^{N-1}$ determines a tridiagonal Hamiltonian matrix of order N in any orthogonal basis $\chi_n^\ell = \sum_{m=0}^{N-1} P_{n,m}^\ell \varphi_m^\ell$ where $\|P_{n,m}^\ell\|$ is an arbitrary orthogonal $(N \times N)$ -matrix of the form

$$\|P_{n,m}^\ell\| = \begin{pmatrix} P_{0,0}^\ell & \cdots & P_{0,N-2}^\ell & 0 \\ \vdots & \vdots & \vdots & \vdots \\ P_{N-2,0}^\ell & \cdots & P_{N-2,N-2}^\ell & 0 \\ 0 & \cdots & 0 & 1 \end{pmatrix}. \quad (59)$$

Let us assume that $\|P_{n,m}^\ell\|$ is the orthogonal transformation matrix that performs the change from $\{\varphi_n^\ell\}_{n=0}^{N-1}$ to the new basis $\{\chi_n^\ell\}_{n=0}^{N-1}$ in which the kinetic energy operator truncated matrix is tridiagonal. To perfect the analogy with the oscillator basis case, let us denote the kinetic energy operator $\frac{2\mu}{\hbar^2} H^0$ (7) tridiagonal matrix in the basis $\{\chi_n^\ell\}_{n=0}^{N-1}$ by $\|T_{n,m}^\ell\|$. The sought for Hamiltonian $\frac{2\mu}{\hbar^2} H$ matrix $\|h_{n,m}\|$ of order N is presumed to be of a Jacobi form (6) in the basis $\{\chi_n^\ell\}_{n=0}^{N-1}$.

Thus, the first $N-1$ of the wave function $\psi(k, r)$ expansion coefficients in the combined basis set $\{\{\chi_n^\ell\}_{n=0}^{N-1}, \{\phi_n^\ell\}_{n=N}^\infty\}$ obey the equations

$$\begin{aligned} a_0 c_0(k) + b_1 c_1(k) &= k^2 c_0(k) \\ b_n c_{n-1}(k) + a_n c_n(k) + b_{n+1} c_{n+1}(k) &= k^2 c_n(k), \quad n = 1, \dots, N-2. \end{aligned} \quad (60)$$

It is easy to check that a sufficient condition that the algebraic version of the Marchenko method be applicable for the construction of the tridiagonal Hamiltonian matrix (6) is that

$$a_{n+1} = T_{n+1,n+1}^\ell, \quad b_{n+1} = T_{n,n+1}^\ell, \quad \text{for } n = M, \dots, N-2, \quad M = \lceil \frac{N}{2} \rceil. \quad (61)$$

If N is odd, to (61) must be added the constraint that $a_M = T_{M,M}^\ell$. In this case $c_{M+1} = f_{M+1}$, $c_M = f_M$, where f_n satisfy the “free” equations

$$T_{n,n-1}^\ell f_{n-1}(k) + T_{n,n}^\ell f_n(k) + T_{n,n+1}^\ell f_{n+1}(k) = k^2 f_n(k), \quad n = 1, \dots, N-2, \quad (62)$$

and we obtain for $n \leq M-1$

$$c_n(k) = \sum_{m=n}^{N-n-1} K_{n,m} f_m(k). \quad (63)$$

Notice that in going from the initial Laguerre basis $\{\phi_n^\ell\}_{n=0}^\infty$ to the combined basis set $\{\{\chi_n^\ell\}_{n=0}^{N-1}, \{\phi_n^\ell\}_{n=N}^\infty\}$ the submatrices $\|h_{n,m}^0\|_{n,m=0}^{N-1}$ and $\|A_{n,m}^\ell\|_{n,m=0}^{N-1}$ are transformed into $\|T_{n,m}^\ell\|_{n,m=0}^{N-1}$ and the identity matrix of order N respectively. In addition, the elements $h_{N-1,N}^0$, $A_{N-1,N}^\ell$ and $h_{N,N-1}^0$, $A_{N,N-1}^\ell$ are multiplied by d_{N-1}^ℓ . The rest of the (infinite) matrices $\|h_{n,m}^0\|$ and $\|A_{n,m}^\ell\|$ is unaltered. It thus follows that f_n must satisfy (in addition to (62)) the equations

$$T_{N-1,N-2}^\ell f_{N-2}(k) + T_{N-1,N-1}^\ell f_{N-1}(k) + d_{N-1}^\ell J_{N-1,N}^\ell f_N(k) = k^2 f_{N-1}(k), \quad (64)$$

$$d_{N-1}^\ell J_{N,N-1}^\ell(k) f_{N-1}(k) + J_{N,N}^\ell(k) f_N(k) + J_{N,N+1}^\ell(k) f_{N+1}(k) = 0. \quad (65)$$

$$J_{n,m}^\ell(k) f_m(k) = 0, \quad n = N+1, \dots. \quad (66)$$

Putting $f_n = \mathcal{S}_{n,\ell}$ for $n \geq N$ and $f_{N-1} = \mathcal{S}_{N-1,\ell}/d_{N-1}^\ell$ [in view of the equation (66) and (65), respectively], from Eq. (64) by using the tree-term recursion relation (62) we obtain the coefficients $\tilde{\mathcal{S}}_{n,\ell}$ with $n = 0, \dots, N-2$. Similarly, setting $f_n = \mathcal{C}_{n,\ell}^{(\pm)}$ for $n \geq N$ and $f_{N-1} = \mathcal{C}_{N-1,\ell}^{(\pm)}/d_{N-1}^\ell$, we obtain the coefficients $\tilde{\mathcal{C}}_{n,\ell}^{(\pm)}$ with $n = 0, \dots, N-2$. From the Wronskian-like relation (see e.g. [16])

$$J_{n+1,n}^\ell(k) \left(\mathcal{C}_{n+1,\ell}^{(\pm)}(k) \mathcal{S}_{n,\ell}(k) - \mathcal{C}_{n,\ell}^{(\pm)}(k) \mathcal{S}_{n+1,\ell}(k) \right) = k, \quad n \geq 0 \quad (67)$$

it follows that

$$T_{n+1,n}^\ell \left(\tilde{\mathcal{C}}_{n+1,\ell}^{(\pm)}(k) \tilde{\mathcal{S}}_{n,\ell}(k) - \tilde{\mathcal{C}}_{n,\ell}^{(\pm)}(k) \tilde{\mathcal{S}}_{n+1,\ell}(k) \right) = k, \quad 0 \leq n \leq N-2. \quad (68)$$

Besides, since the system of equations (52) in $\mathcal{S}_{n,\ell}$ is homogeneous, the sets $\{\mathcal{S}_{n,\ell}\}_{n=0}^\infty$ and $\{\tilde{\mathcal{S}}_{n,\ell}\}_{n=0}^\infty$ are connected by a linear transformation and therefore $\tilde{\mathcal{S}}_{n,\ell}$ also satisfy the homogeneous equation

$$T_{0,0}^\ell \tilde{\mathcal{S}}_{0,\ell}(k) + T_{0,1}^\ell \tilde{\mathcal{S}}_{1,\ell}(k) = k^2 \tilde{\mathcal{S}}_{0,\ell}(k), \quad (69)$$

whereas $\tilde{\mathcal{C}}_{n,\ell}^{(\pm)}$ obey the inhomogeneous one

$$T_{0,0}^\ell \tilde{\mathcal{C}}_{0,\ell}^{(\pm)}(k) + T_{0,1}^\ell \tilde{\mathcal{C}}_{1,\ell}^{(\pm)}(k) = k^2 \tilde{\mathcal{C}}_{0,\ell}^{(\pm)}(k) + \frac{k}{\tilde{\mathcal{S}}_{0,\ell}(k)}. \quad (70)$$

Thus, the two sets, $\{\tilde{\mathcal{S}}_{n,\ell}\}_{n=0}^\infty$ and $\{\tilde{\mathcal{C}}_{n,\ell}\}_{n=0}^\infty$, $\tilde{\mathcal{C}}_{n,\ell} = \frac{1}{2}(\tilde{\mathcal{C}}_{n,\ell}^{(-)} + \tilde{\mathcal{C}}_{n,\ell}^{(+)})$, are “free” independent respectively sine-like [$\mathcal{S}_{n,\ell} = \mathcal{S}_{n,\ell}$, $n \geq N$] and cosine-like [$\tilde{\mathcal{C}}_{n,\ell}^{(\pm)} = \mathcal{C}_{n,\ell}^{(\pm)}$, $n \geq N$] solutions to Eqs. (62)-(66) (see e.g. [7]).

From the above discussion it follows that to obtain f_n with $0 \leq n \leq N-1$, which are involved in Eq. (63), we can place $f_N = w_N$, $f_{N-1} = w_{N-1}/d_{N-1}^\ell$, where w_n are defined by (49). Then, inserting this f_N and f_{N-1} in Eq. (64) gives f_{N-2} . Once f_{N-2} and f_{N-1} are known, f_n for $n = N-3, \dots, 0$ are obtained by using the tree-term recursion relation (62). $K_{n,m}$ are determined by the equations (27) and (28) [in which the upper limit in the sums is equal to $N-n-1$]. The expressions for $\{a_n\}$, $\{b_n\}$ are the same as (30).

$Q_{n,n'}$ with $n \leq N-1$ in Eqs. (27) and (28) are defined [in view of the overlap matrix form in the combined basis] by

$$Q_{n,n'} = \frac{2}{\pi} \int_0^\infty f_n(k) \overline{f_{n'}(k)} + \sum_\nu f_n(i\kappa_\nu) \overline{f_{n'}(i\kappa_\nu)}, \quad n' \leq N-2, \quad (71)$$

$$\begin{aligned} Q_{n,N-1} = & \frac{2}{\pi} \int_0^\infty f_n(k) [\overline{w_{N-1}(k)}/d_{N-1}^\ell + d_{N-1}^\ell A_{N-1,N}^\ell \overline{w_N(k)}] + \\ & + \sum_\nu f_n(i\kappa_\nu) [\overline{w_{N-1}(i\kappa_\nu)}/d_{N-1}^\ell + d_{N-1}^\ell A_{N-1,N}^\ell \overline{w_N(i\kappa_\nu)}], \end{aligned} \quad (72)$$

$$Q_{n,n'} = \frac{2}{\pi} \int_0^\infty f_n(k) A_{n',m}^\ell \overline{w_m(k)} + \sum_\nu f_n(i\kappa_\nu) A_{n',m}^\ell \overline{w_m(i\kappa_\nu)}, \quad n' \geq N. \quad (73)$$

Notice that at large k , as seen in Eq. (50), $\mathcal{C}_{n,\ell}^{(\pm)}(k) \sim k^\ell$. Thus, as in the case of the oscillator basis, we should restrict ourselves to the description of the scattering data on a finite energy interval, beyond the boundary of which the phase shift needs generally to be modified to provide the convergence of the integrals in Eqs. (71)-(73).

6 Conclusion

In the potential scattering case the finite-difference approach and J-matrix method share the tridiagonal representation of the Hamiltonian. The analogy can be carried over to the inverse scattering problem formalism. Here, the

J-matrix version of the Marchenko equation algebraic analogue is formulated and its numerical realization features are considered. The merit of JME is that it is free from a parameter fit inherent in the previous J-matrix inverse scattering approach [2, 3, 4, 5]. We also construct a tridiagonal Hamiltonian matrix of some order M in an orthogonalized Laguerre basis; in doing so it is sufficient to tridiagonalize the matrix representation of the reference Hamiltonian H^0 in the finite orthogonal basis subset of size $N = 2M$. As has been shown in Ref. [5], in the two coupled-channel case without threshold the sought-for interaction matrix may be of a “quasi-tridiagonal” form. On this assumption JME can be easily expanded to multichannel scattering.

Acknowledgments

The author acknowledge helpful conversations with A .M. Shirokov. This work was partially supported by the State Program “Russian Universities”, by the Russian Foundation of Basic Research grant No 02-02-17316.

References

- [1] E. J. Heller, H. A. Yamani, Phys. Rev. A **9**, 1209 (1974).
- [2] S. A. Zaitsev, Teoret. Mat. Fiz. **115**, 263 (1998) [Theor. Math. Phys. **115**, 575 (1998)].
- [3] S. A. Zaitsev, Teoret. Mat. Fiz. **121**, 424 (1999) [Theor. Math. Phys. **121**, 1617 (1999)].
- [4] S. A. Zaitsev, E. I. Kramar, J. Phys. G, **27**, 2037 (2001).
- [5] A. M. Shirokov, A. I. Mazur, S. A. Zaytsev, J. P. Vary, T. A. Weber, Phys. Rev. C **70** (2004) 044005.
- [6] H. A. Yamani, A. D. Alhaidari, and M. S. Abdelmonem, Phys. Rev. A **64**, 042703 (2001).
- [7] H. A. Yamani, L. Fishman, J. Math. Phys. **16**, 410 (1975).
- [8] K. Ghanbari, IP **17**, 211 (2001).
- [9] G. M. L. Gladwell, N. B. Willms, IP **5**, 165 (1989).
- [10] V. M. Chabanov, J. Phys. A, **37**, 9139 (2004).
- [11] K. M. Case, J. Math. Phys. **14**, 916 (1973)

- [12] B. N. Zakhariev, A. A. Suzko, *Direct and inverse problems. In: Potentials in quantum scattering* (2-nd ed. Berlin, Heidelberg, New York: Springer-Verlag, 1990).
- [13] A. I. Baz, Ya. B. Zeldovitch, and A. M. Perelomov, *Scattering, Reactions and Decays in Non-relativistic Quantum Mechanics* (Moscow: Nauka, 1971).
- [14] K. Chadan, P. C. Sabatier, *Inverse Problems in Quantum Scattering Theory* (New York, Heidelberg, Berlin: Springer-Verlag, 1977).
- [15] J. T. Broad, Phys. Rev. A **31**, 1494 (1985).
- [16] J. T. Broad, W. P. Reinhardt, J. Phys. B **9**, 1491 (1976).




Figure 1:




Figure 2:




Figure 3:

Table

$N = 7, \rho = \frac{R}{2}$				
$\kappa R = .6512647458,$ $\mathcal{M}R^{1/2} = 1.6017576599$			$\kappa R = .6380449999,$ $\mathcal{M}R^{1/2} = 1.5833238674$	
j	$Z_{N-1,j}$	λ_j	$Z_{N-1,j}$	λ_j
0	0.0356514517	-0.0381260178	0.0362075259	-0.0353279575
1	0.1482712147	0.4605384282	0.1482712147	0.4605384282
2	0.2309801539	1.3246452781	0.2309801539	1.3246452781
3	0.3094585382	2.5044702689	0.3094585382	2.5044702689
4	0.4084394267	4.1865934000	0.4084394267	4.1865934000
5	0.5275945630	6.7063360348	0.5275945630	6.7063360348
6	0.6184249465	10.1425219887	0.6183926387	10.1425219887